

HRP-conjugated 6His, His-Tag Monoclonal antibody

Cat: B30001T-HRP

Company: HaoKebio

Uniprot ID: HISTAG

Applications:


WB:1:5000-1:50000

Organism: Mouse / IgG1

Species reactivity: Recombinant protein

Background:

Protein tags are a protein or peptide sequences located either on the C- or N- terminal of the target protein. His-tag is often used for affinity purification and binding assays. Expressed. The His-tag antibody is a useful tool for monitoring of the His-tagged proteins and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells. A His-tag (polyhistidine tag) consists of at least six histidine residues that are located at the N- or C-terminus of recombinant proteins. It is commonly used for affinity purification and protein binding experiments.



1.Multi-tag Protein
 2.Multi-tag Protein in 293
 3.Multi-tag Protein in HeLa

Dilution of 1:50000 incubated at room temperature for 1.5 hours.

Source of Reagents:

发表[中文论文]请标注:HRP-conjugated 6His, His-Tag(B30001T-HRP)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:HRP-conjugated 6His, His-Tag(B30001T-HRP) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

Protein full name:

6*His, His-Tag

Synonyms:

6*His, His Tag, His Tag, 6*His, 6*his tag, 6*his, His tag

Immunogen:

Recombinant protein

Isotype:

IgG1

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Images: